


This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1.(Original) Salts of saturated, partially or fully unsaturated, heterocyclic cations having the bis(trifluoromethyl)imide anion, $N(CF_3)_2^-$, which have the general formula (1)



where

X = N, P, O or S

n = an integer selected from 0, 1 or 2 in such a way that X is saturated in accordance with its valency increased by 1,

A = a saturated, partially or fully unsaturated 3- to 8-membered hydrocarbon chain,
in which all carbon atoms apart from one may be replaced by identical or different heteroatoms selected from N, P, O and S,
where the carbon atoms of the hydrocarbon chain and the heteroatoms present therein are saturated by substituents R^2 in accordance with their valency,

$R^1, R^2 =$ -H, with the proviso that there is no bond to the positively charged hetero-atom,
straight-chain or branched alkyl having 1-20 carbon atoms
straight-chain or branched alkenyl having 2-20 carbon atoms and one or more double bonds
straight-chain or branched alkynyl having 2-20 carbon atoms and one or more triple bonds
saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms

halogen, in particular fluorine or chlorine, with the proviso that, for X = N, O, S, there is no halogen-heteroatom bond,
 -NO₂, with the proviso that there is no bond to a positively charged heteroatom,
 -CN, with the proviso that there is no bond to a positively charged heteroatom,

where the R² and/or R¹ in different and/or identical position of the heterocyclic ring are in each case identical or different,

where the R² and/or R¹ may be connected to one another in pairs by a single or double bond,

where one or more R² and/or R¹ may be partially or fully substituted by halogens, in particular -F and/or -Cl, or partially by -CN or -NO₂, with the proviso that not all R² and R¹ are fully halogenated,

and where one or two carbon atoms of the R¹ and/or R² may be replaced by heteroatoms and/or atomic groups selected from the group -O-, -C(O)-, C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅, where the α-position of the R¹ is not replaced for X = O, S.

2. (Original) Salts according to Claim 1, in which A is a 4-, 5- or 6-membered hydrocarbon chain.

3. (Currently Amended) Salts according to Claim 1 or 2, in which A is a hydrocarbon chain in which zero, one or two carbon atoms are replaced by heteroatoms selected from N, P, O and S.


4. (Currently Amended) Salts according to ~~one of Claims 1 to 3~~ Claim 1, in which R¹, R², independently of one another, have the meaning

-H, with the proviso that there is no bond to the positively charged heteroatom, halogen, in particular fluorine, with the proviso that for X = N, O, S, there is no halogen-heteroatom bond,

straight-chain or branched alkyl having 1-6 carbon atoms, in particular -CH₃, -C₂H₅, -*n*-C₃H₇, -CH(CH₃)₂, -*n*-C₄H₉, -*n*-C₆H₁₃,

straight-chain or branched, partially or perfluorinated alkyl having 1-6 carbon atoms, in particular -CF₃, -C₂F₅, -C₄F₉.

5. (Currently Amended) Salts according to ~~one of Claims 1 to 4~~ Claim 1, in which the saturated, partially or fully unsaturated, heterocyclic cation is selected from the following group:



where

X = N, P, O or S


Y = in each case, independently of one another, N, P, O, S or C,
where at least one Y = C

n = 0 for unsaturated X = O, S
1 for saturated X = O, S or for unsaturated X = N, P
2 for saturated X = N, P

m = 0 for saturated Y = O, S or for unsaturated Y = N, P
1 for saturated Y = N, P or for Y = sp²-C
2 for Y = sp³-C

where the radicals R¹, R² are as defined in Claim 1.

6. (Currently Amended) Salts according to ~~one of Claims 1 to 4~~ Claim 1, in which the saturated, partially or fully unsaturated, heterocyclic cation is selected from the following group:



where

X = in each case, independently of one another, N, P, O, S or C,
where at least one X = N, P, O or S


n = 0 for saturated X = O, S or for unsaturated X = N, P
1 for saturated X = N, P or for X = sp^2 -C
or for substitution on the sp^2 ring carbon atom
2 for X = sp^3 -C
or for substitution on the sp^3 ring carbon atom

where n is increased by 1 for one X = N, P, O or S

A' = saturated, partially or fully unsaturated 2- to 7-membered hydrocarbon chain,
where the carbon atoms of the hydrocarbon chain are saturated by
substituents R^2 in accordance with their valency

where the radicals R^1 , R^2 are as defined in Claim 1, with the proviso that the
substituents R^1 in the α -position to the positively charged heteroatom do not have a
methylene group directly adjacent to the heterocyclic ring.

7. (Currently Amended) Salts according to ~~one of Claims 1 to 4~~ Claim 1, in which the
saturated, partially or fully unsaturated, heterocyclic cation has the following structure



where

X = N or P

Y = N, P, O or S

n = 0 for Y = O, S
1 for Y = N, P


A' = saturated, partially or fully unsaturated 2- to 7-membered hydrocarbon chain,

B' = saturated, partially or fully unsaturated 1- to 6-membered hydrocarbon chain,

where all carbon atoms of the hydrocarbon chains A' and B' apart from one may be replaced by identical or different heteroatoms selected from N, P, O and S
and where the carbon atoms of the hydrocarbon chains A' and B' are saturated by substituents R² in accordance with their valency

and where the radicals R¹, R² are as defined in Claim 1.

8. (Original) Salts of saturated, partially or fully unsaturated, heterocyclic di-cations having the bis(trifluoromethyl)imide anion, N(CF₃)₂⁻, which have the general formula (2)



where

X, Y = each, independently of one another, N, P, O or S

$n, m =$ an integer selected from 0, 1 or 2 in such a way that X and Y are each saturated in accordance with their valency increased by 1,

$A'', B'' =$ saturated, partially or fully unsaturated 0- to 4-membered hydrocarbon chain,
in which the carbon atoms may be replaced by identical or different heteroatoms selected from N, P, O and S,
where at least one carbon atom is present in the chains A'' and B'' together and
where the carbon atoms of the hydrocarbon chains A'' and B'' and the heteroatoms present therein are saturated by substituents R^2 in accordance with their valency,

$R^1, R^2, R^3 =$ -H, with the proviso that there is no bond to the positively charged heteroatom,
straight-chain or branched alkyl having 1-20 carbon atoms
straight-chain or branched alkenyl having 2-20 carbon atoms and one or more double bonds
straight-chain or branched alkynyl having 2-20 carbon atoms and one or more triple bonds
saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms
halogen, in particular fluorine or chlorine, with the proviso that, for X, = N, O, S, there is no halogen-heteroatom bond,
-NO₂, with the proviso that there is no bond to a positively charged heteroatom,
-CN, with the proviso that there is no bond to a positively charged heteroatom,

where the R^1, R^2 and/or R^3 in different and/or identical position of the heterocyclic ring are in each case identical or different,

where the R^1, R^2 and/or R^3 may be connected to one another in pairs by a

single or double bond,


where one or more R^1 , R^2 and/or R^3 may be partially or fully substituted by halogens, in particular -F and/or -Cl, or partially by -CN or -NO₂, with the proviso that that not all R^1 , R^2 and R^3 are fully halogenated,

and where one or two carbon atoms of the R^1 , R^2 and/or R^3 may be replaced by heteroatoms and/or atomic groups selected from the group -O-, -C(O)-, C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅, where the α -positions of the R^1 and of the R^3 are not replaced for X = O, S or Y = O, S,

where the heterocyclic di-cation is a 4-, 5-, 6-, 7-, 8- or 9-membered ring.

9. (Original) Salts according to Claim 8, in which the heterocyclic di-cation is a 5-, 6- or 7-membered ring.

10. (Original) Salts of saturated, partially or fully unsaturated, heterocyclic cations having the bis(trifluoromethyl)imide anion, $N(CF_3)_2^-$, which have the general formula (3)



where

X, Y = each, independently of one another, N, P, O or S

n = an integer selected from 0, 1 or 2 in such a way that X is saturated in accordance with its valency increased by 1,

m = an integer selected from 0, 1 or 2 in such a way that Y is saturated in

accordance with its valency,

$R^1, R^2 =$ -H, with the proviso that there is no bond to the positively charged heteroatom,
straight-chain or branched alkyl having 1-20 carbon atoms
straight-chain or branched alkenyl having 2-20 carbon atoms and one or more double bonds
straight-chain or branched alkynyl having 2-20 carbon atoms and one or more triple bonds
saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms
halogen, in particular chlorine, with the proviso that, for X, = N, O, S, there is no halogen-heteroatom bond,
-OR, with the proviso that the substituted heteroatom is not O or S,


where the R^2 and/or R^1 in different and/or identical position of the heterocyclic ring are in each case identical or different,

where the R^2 and/or R^1 may be connected to one another in pairs by a single or double bond,

where one or more R^2 and/or R^1 may be partially or fully substituted by halogens, in particular -F and/or -Cl, or partially by -CN or -NO₂, with the proviso that not all R^2 and R^1 are fully halogenated,


and where one or two carbon atoms of the R^1 and/or R^2 may be replaced by heteroatoms and/or atomic groups selected from the group -O-, -C(O)-, C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅, where the α -position of the R^1 is not replaced for X = O, S.

11. (Currently Amended) Process for the preparation of salts according to ~~one of Claims 1 to 7~~ Claim 1, characterised in that an alkali metal fluoride of the general formula DF, where D selected from the group of the alkali metals, is reacted in a polar organic solvent with



where $R^F = F$ or C_pF_{2p+1} , where $p = 1 - 8$,

and a salt of the general formula (4)



where

$E^- = F^-, Cl^-, Br^-, I^-, BF_4^-, ClO_4^-, AsF_6^-, SbF_6^-, SbCl_6^-, PF_6^-, R^F SO_3^-, FSO_3^-,$
 $(R^F)_2P(O)O^-, R^F P(O)_2O^-, RSO_3^-, ROSO_3^-, \frac{1}{2}SO_3^{2-}, CN^-, SCN^-, R^F C(O)O^-,$
 $RC(O)O^-,$ 2,4-dinitrophenolate or 2,4,6-trinitrophenolate, where R^F is a
 perfluorinated C_1 to C_8 -alkyl group or perfluorinated aryl group and R is a
 C_1 to C_8 -alkyl group or aryl group

$X =$ N, P, O or S

$n =$ an integer selected from 0, 1 or 2 in such a way that X is saturated in
 accordance with its valency increased by 1,

$A =$ a saturated, partially or fully unsaturated 3- to 8-membered hydrocarbon
 chain,
 in which all carbon atoms apart from one may be replaced by identical or
 different heteroatoms selected from N, P, O and S,
 where the carbon atoms of the hydrocarbon chain and the heteroatoms
 present therein are saturated by substituents R^2 in accordance with their
 valency,

$R^1, R^2 =$ -H, with the proviso that there is no bond to the positively charged hetero-
 atom,
 straight-chain or branched alkyl having 1-20 carbon atoms
 straight-chain or branched alkenyl having 2-20 carbon atoms and one or
 more double bonds
 straight-chain or branched alkynyl having 2-20 carbon atoms and one or

more triple bonds
saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms
halogen, in particular fluorine or chlorine, with the proviso that, for X = N,
O, S, there is no halogen-heteroatom bond,
-NO₂, with the proviso that there is no bond to a positively charged hetero-
atom,
-CN, with the proviso that there is no bond to a positively charged hetero-
atom,


where the R² and/or R¹ in different and/or identical position of the
heterocyclic ring are in each case identical or different,

where the R² and/or R¹ may be connected to one another in pairs by a
single or double bond,

where one or more R² and/or R¹ may be partially or fully substituted by
halogens, in particular -F and/or -Cl, or partially by -CN or -NO₂, with the
proviso that not all R² and R¹ are fully halogenated,


and where one or two carbon atoms of the R¹ and/or R² may be replaced by
heteroatoms and/or atomic groups selected from the group -O-, -C(O)-,
C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and
-PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅,
where the α-position of the R¹ is not replaced for X = O, S.

12. (Currently Amended) Process for the preparation of salts according to ~~one of~~
~~Claims 8 or 9~~ Claim 8, characterised in that an alkali metal fluoride of the general
formula DF, where D selected from the group of the alkali metals, is reacted in a polar
organic solvent with



where R^F = F or C_pF_{2p+1}, where p = 1 – 8,

and a salt of the general formula (5)



where

$E^- = F^-, Cl^-, Br^-, I^-, BF_4^-, ClO_4^-, AsF_6^-, SbF_6^-, SbCl_6^-, PF_6^-, R^F SO_3^-, FSO_3^-, (R^F)_2 P(O)O^-, R^F P(O)_2 O^-, RSO_3^-, ROSO_3^-, \frac{1}{2} SO_3^{2-}, CN^-, SCN^-, R^F C(O)O^-, RC(O)O^-, 2,4\text{-dinitrophenolate or } 2,4,6\text{-trinitrophenolate,}$
 where R^F is a perfluorinated C_1 to C_8 -alkyl group or perfluorinated aryl group and R is a C_1 to C_8 -alkyl group or aryl group

$X, Y =$ each, independently of one another, N, P, O or S

$n, m =$ an integer selected from 0, 1 or 2 in such a way that X, Y are each saturated in accordance with their valency increased by 1,

$A'', B'' =$ saturated, partially or fully unsaturated 0- to 4-membered hydrocarbon chain,
 in which the carbon atoms may be replaced by identical or different heteroatoms selected from N, P, O and S,
 where at least one carbon atom is present in the chains A'' and B'' together and
 where the carbon atoms of the hydrocarbon chains A'' and B'' and the heteroatoms present therein are saturated by substituents R^2 in accordance with their valency,

$R^1, R^2, R^3 =$ -H, with the proviso that there is no bond to the positively charged heteroatom,
 straight-chain or branched alkyl having 1-20 carbon atoms
 straight-chain or branched alkenyl having 2-20 carbon atoms and one or more double bonds
 straight-chain or branched alkynyl having 2-20 carbon atoms and one or

more triple bonds
 saturated, partially or fully unsaturated cycloalkyl having 3-7 carbon atoms
 halogen, in particular fluorine or chlorine, with the proviso that, for X, = N, O, S, there is no halogen-heteroatom bond,
 halogen, in particular fluorine or chlorine, with the proviso that there is no halogen-heteroatom bond,
 -NO₂, with the proviso that there is no bond to a positively charged heteroatom,
 -CN, with the proviso that there is no bond to a positively charged heteroatom,

where the R¹, R² and/or R³ in different and/or identical position of the heterocyclic ring are in each case identical or different,

where the R¹, R² and/or R³ may be connected to one another in pairs by a single or double bond,

where one or more R¹, R² and/or R³ may be partially or fully substituted by halogens, in particular -F and/or -Cl, or partially by -CN or -NO₂, with the proviso that not all R¹, R² and R³ are fully halogenated,

and where one or two carbon atoms of the R¹, R² and/or R³ may be replaced by heteroatoms and/or atomic groups selected from the group -O-, -C(O)-, C(O)O-, -S-, -S(O)-, -SO₂-, -SO₂O-, -N=, -P=, -NH-, -PH-, -NR'- and -PR'- where R' = non-, partially or perfluorinated C₁- to C₆-alkyl or -C₆F₅, where the α-position of the R¹ is not replaced for X = O, S,

where the heterocyclic di-cation is a 4-, 5-, 6-, 7-, 8- or 9-membered ring.

13. (Currently Amended) Process according to Claim 11 ~~or 12~~, characterised in that the alkali metal fluoride employed is KF or RbF.

14. (Currently Amended) Process according to ~~one of Claims 11 to 13~~ Claim 11, characterised in that the reaction takes place at temperatures between -40°C and 80°C, in particular at 0°C to 40°C.

15. (Currently Amended) Process according to ~~one of Claims 11 to 14~~ Claim 11, characterised in that the reaction takes place in a polar organic solvent selected from the group acetonitrile, dimethoxyethane, dimethylformamide and propionitrile.
16. (Currently Amended) Process according to ~~one of Claims 11 to 15~~ Claim 11, characterised in that the reaction is carried out as a one-pot reaction.
17. (Currently Amended) Process according to ~~one of Claims 11 to 16~~ Claim 11, characterised in that the reaction for $E^- = F^-$ is carried out without added alkali metal fluoride DF.
18. (Currently Amended) Process according to ~~one of Claims 11 to 17~~ Claim 11, characterised in that the starting materials for the reaction are employed in approximately equimolar ratio to one another.
19. (Currently Amended) Use of a salt according to ~~one of Claims 1 to 10~~ Claim 1 as ionic liquid.
20. (Currently Amended) Use of a salt according to ~~one of Claims 1 to 10~~ Claim 1 as non-aqueous electrolyte.
21. (Currently Amended) Use of a salt according to ~~one of Claims 1 to 10~~ Claim 1 as reagent for the introduction of $N(CF_3)_2$ groups.
22. (Currently Amended) Use of a salt according to ~~one of Claims 1 to 10~~ Claim 1 as phase-transfer catalyst.
23. (Currently Amended) Use of a salt according to ~~one of Claims 1 to 10~~ Claim 1 as intermediate for the synthesis of liquid-crystal compounds or active ingredients, in particular for medicaments or crop-protection agents.